arXiv:1610.09794v1 [math.DG] 31 Oct 2016

WILLMORE INEQUALITY ON HYPERSURFACES IN HYPERBOLIC SPACE

YINGXIANG HU

Abstract. In this article, we prove a geometric inequality for star-shaped and mean-convex hypersurfaces in hyperbolic space by inverse mean curvature flow. This inequality can be considered as a generalization of Willmore inequality for closed surface in hyperbolic 3-space.

1. Introduction

The classical isoperimetric inequality and its generalization, the Alexandrov-Fenchel inequalities play an important role in different branches of geometry. Let $\Omega \subset \mathbb{R}^n$ be a smooth bounded domain with boundary Σ, then the classical isoperimetric inequality is

$$|\Sigma| \geq n^{\frac{n-1}{n}} \omega_{n-1}^{\frac{1}{n}} |\Omega|^{\frac{n-1}{n}}, \tag{1}$$

and equality in (1) holds if and only if Ω is a geodesic ball.

For $k \in \{1, \cdots, n-1\}$, we denote by p_k the normalized k-th order mean curvature of Σ, and set $p_0 = 1$ by convention. The celebrated Alexandrov-Fenchel inequalities [1, 2, 15] for convex hypersurface $\Sigma^{n-1} \subset \mathbb{R}^n$ are

$$\frac{1}{\omega_{n-1}} \int_\Sigma p_k d\mu \geq \left(\omega_{n-1} \int_\Sigma p_j d\mu \right)^{\frac{n-1-k}{n-1-j}}, \quad 0 \leq j < k \leq n-1, \tag{2}$$

and equality in (2) holds if and only if Ω is a geodesic ball.

Observe that the isoperimetric inequality holds for non-convex domains, it is natural to extend the original Alexandrov-Fenchel inequality to non-convex domains, see [20, 31, 21, 7, 8, 9, 28]. We should also mention that the Willmore inequality, which is a weaker form of Alexandrov-Fenchel inequality, has been established for closed surfaces in $\mathbb{R}^3$. More precisely, for any closed surface $\Sigma \subset \mathbb{R}^3$, the Willmore inequality [10, 26, 30] is

$$\int_\Sigma p_1^2 d\mu \geq \omega_2, \tag{3}$$

and equality in (3) holds if and only if Σ is a geodesic sphere.

It is interesting to establish the Alexandrov-Fenchel inequalities for hypersurfaces in hyperbolic space, see [4, 16]. Recently, the following hyperbolic Alexandrov-Fenchel inequalities were obtained.

Theorem A ([17, 18, 32]). *Let $k \in \{1, \cdots, n-1\}$. Any horospherical convex hypersurface $\Sigma \subset \mathbb{H}^n$ satisfies*

$$\int_\Sigma p_k d\mu \geq \omega_{n-1} \left[\left(\frac{|\Sigma|}{\omega_{n-1}} \right)^{\frac{2}{k}} + \left(\frac{|\Sigma|}{\omega_{n-1}} \right)^{\frac{2(n-1-k)}{k(n-1)}} \right]^{\frac{k}{2}}. \tag{4}$$

Equality in (4) holds if and only if Σ is a geodesic sphere.

Inequality (4) was proved in [17] for $k = 4$ and in [18] for general even k. For $k = 1$, (4) was proved in [18] with a help of a result of Cheng and Zhou [12]. For general integer k, (4) was proved in [32].

For $k = 2$, inequality (4) was proved by Li-Wei-Xiong [25] under a weaker condition that Σ is star-shaped and two-convex. More precisely,

Theorem B ([25]). *Any star-shaped and 2-convex hypersurface $\Sigma \subset \mathbb{H}^n (n \geq 3)$ satisfies*

$$\int_\Sigma p_2 d\mu \geq \omega_{n-1}^{\frac{2}{n-1}} |\Sigma|^{\frac{n-3}{n-1}} + |\Sigma|. \tag{5}$$

Equality in (5) holds if and only if Σ is a geodesic sphere.

The Willmore inequality (3) has also been generalized to closed surface $\Sigma \subset \mathbb{H}^3$.

Theorem C ([11, 27, 29]). *Any closed surface $\Sigma \subset \mathbb{H}^3$ satisfies*

$$\int_\Sigma p_1^2 d\mu \geq \omega_2 + |\Sigma|. \tag{6}$$

Equality in (6) holds if and only if Σ is a geodesic sphere.

Inspired by these previous results, we prove the following geometric inequality, which can be considered as a generalization of Willmore inequality for hypersurfaces in hyperbolic space.

Theorem 1. *Let $\Sigma \subset \mathbb{H}^n$ $(n \geq 3)$ be star-shaped and mean-convex hypersurface, then*

$$\int_\Sigma p_1^2 d\mu \geq \omega_{n-1}^{\frac{2}{n-1}} |\Sigma|^{\frac{n-3}{n-1}} + |\Sigma|. \tag{7}$$

Equality in (7) holds if and only if Σ is a geodesic sphere.

We expect that the inequality (7) will be useful in defining the Hawking mass for hypersurfaces in $\mathbb{H}^n$. In [24], the *Hawking mass* for a closed embedded surface Σ in $\mathbb{H}^3$ is defined as

$$m_H(\Sigma) = \frac{|\Sigma|^{\frac{1}{2}}}{2\omega_2^{\frac{1}{2}}} \left[1 - \omega_2^{-1} \int_\Sigma (p_1^2 - 1) d\mu \right].$$

We now give the outline of the proof of Theorem 1. Motivated by [6, 13, 25], we adopt the inverse mean curvature flow (IMCF) in our proof. This flow has been used by Huisken and Ilmanen [22, 23] to prove the Riemannian Penrose inequality in general relativity. We start from a given star-shaped and mean-convex hypersurface Σ, and evolve it by IMCF. By the convergence results of Gerhardt [19] (see also [6]), the IMCF exists for all time, and the evolving hypersurface Σ_t with $\Sigma_0 = \Sigma$ remains star-shaped and mean-convex for all $t \geq 0$.

We next consider the quantity:

$$Q(t) := |\Sigma_t|^{-\frac{n-3}{n-1}} \int_{\Sigma_t} (p_1^2 - 1) d\mu.$$

We study the limit of $Q(t)$ as $t \to \infty$. Notice that the roundness estimate for Σ_t is not strong enough to calculate the limit of $Q(t)$. However, similar to [6, 25], we are able to give a positive lower bound for the limit of $Q(t)$, which will be used to establish the monotonicity of $Q(t)$. Finally, we prove that $Q(t)$ is monotone non-increasing under IMCF. From this, Theorem 1 follows immediately.

2. Preliminaries

In this article, we consider the hyperbolic space $\mathbb{H}^n = \mathbb{R}^+ \times \mathbb{S}^{n-1}$ equipped with the metric

$$\overline{g} = dr^2 + \sinh^2 r g_{\mathbb{S}^{n-1}},$$

where $g_{\mathbb{S}^{n-1}}$ is the standard round metric on the unit sphere $\mathbb{S}^{n-1}$. Let $\Sigma \subset \mathbb{H}^n$ be a closed hypersurface with its unit outward normal vector ν. The second fundamental form h of Σ is defined by

$$h(X, Y) = \langle \overline{\nabla}_X \nu, Y \rangle$$

for any $X, Y \in T\Sigma$. The principal curvature $\kappa = (\kappa_1, \cdots, \kappa_n)$ are the eigenvalues of h with respect to the induced metric g on Σ. For $k \in \{1, \cdots, n-1\}$, the normalized k-th elementary symmetric polynomial of κ is defined as

$$p_k(\kappa) := \frac{1}{\binom{n-1}{k}} \sum_{i_1 < i_2 < \cdots < i_k} \kappa_{i_1} \cdots \kappa_{i_k},$$

which can also be viewed as a function of the second fundamental form $h_i^j = g^{jk} h_{ki}$. For abbreviation, we write p_k for $p_k(\kappa)$.

We now consider the inverse mean curvature flow (IMCF)

$$\partial_t X = \frac{1}{(n-1)p_1} \nu. \tag{8}$$

where $\Sigma_t = X(t, \cdot)$ is a family of hypersurfaces in $\mathbb{H}^n$, ν is the unit outward normal to $\Sigma_t = X(t, \cdot)$. Let $d\mu_t$ be its area element on Σ_t. We list the following evolution equations.

Lemma 2. *Under IMCF (8), we have:*

$$\partial_t p_1 = -\frac{1}{(n-1)^2} \Delta \left(\frac{1}{p_1} \right) - \frac{1}{(n-1)^2 p_1} (|A|^2 + n - 1). \tag{9}$$

$$\partial_t d\mu = d\mu. \tag{10}$$

In [19], Gerhardt investigated the inverse curvature flow of star-shaped hypersurfaces in hyperbolic space.

Theorem 3 ([19])**.** *If the initial hypersurface is star-shaped and mean-convex, then the solution for IMCF (8) exists for all time t and preserves the condition of star-shapedness and mean-convexity. Moreover, the hypersurfaces become strictly convex exponentially fast and more and more totally umbilical in the sense of*

$$|h_i^j - \delta_i^j| \leq C e^{-\frac{t}{n-1}}, \quad t > 0,$$

i.e., the principal curvatures are uniformly bounded and converge exponentially fast to one.

3. The asymptotic behavior of monotone quantity

We define the quantity

$$Q(t) := |\Sigma_t|^{-\frac{n-3}{n-1}} \int_{\Sigma_t} (p_1^2 - 1) d\mu,$$

where $|\Sigma_t|$ is the area of Σ_t. In this section, we estimate the lower bound of the limit of $Q(t)$. First of all, we recall the following sharp Sobolev inequality on $\mathbb{S}^{n-1}$ due to Beckner [3]. This Sobolev inequality is crucial in analyzing the asymptotic behavior of the monotone quantity, see [6, 25], etc.

Lemma 4. *For every positive function f on $\mathbb{S}^{n-1}$, we have*

$$\begin{aligned} &\int_{\mathbb{S}^{n-1}} f^{n-3} dvol_{\mathbb{S}^{n-1}} + \frac{n-3}{n-1} \int_{\mathbb{S}^{n-1}} f^{n-5} |\nabla f|^2 dvol_{\mathbb{S}^{n-1}} \\ \geq &\omega_{n-1}^{\frac{2}{n-1}} \left(\int_{\mathbb{S}^{n-1}} f^{n-1} dvol_{\mathbb{S}^{n-1}} \right)^{\frac{n-3}{n-1}}. \end{aligned} \tag{11}$$

Moreover, equality in (11) holds if and only if f is a constant.

Proof. From Theorem 4 in [3], for any positive smooth function w on $\mathbb{S}^{n-1}$, we have the following inequality

$$\begin{aligned} &\frac{4}{(n-1)(n-3)} \int_{\mathbb{S}^{n-1}} |\nabla w|^2 dvol_{\mathbb{S}^{n-1}} + \int_{\mathbb{S}^{n-1}} w^2 dvol_{\mathbb{S}^{n-1}} \\ \geq &\omega_{n-1}^{\frac{2}{n-1}} \left(\int_{\mathbb{S}^{n-1}} w^{\frac{2(n-1)}{n-3}} dvol_{\mathbb{S}^{n-1}} \right)^{\frac{n-3}{n-1}}. \end{aligned} \tag{12}$$

Moreover, the equality holds if and only if w is a constant. For any positive function f on $\mathbb{S}^{n-1}$, by letting $w = f^{\frac{n-3}{2}}$, we obtain the desired estimate. □

Proposition 5. *Under IMCF (8), we have*

$$\liminf_{t \to \infty} Q(t) \geq \omega_{n-1}^{\frac{2}{n-1}}. \tag{13}$$

Proof. Recall that star-shaped hypersurfaces can be written as graphs of function $r = r(t, \theta)$, $\theta \in \mathbb{S}^{n-1}$. Denote $\lambda(r) = \sinh(r)$, then $\lambda'(r) = \cosh(r)$. We next define a function $\varphi(\theta) = \Phi(r(\theta))$, where $\Phi(r)$ is a positive function satisfying $\Phi' = \frac{1}{\lambda}$. Let $\theta = \{\theta^j\}$, $j = 1, \cdots, n-1$ be a coordinate system on $\mathbb{S}^{n-1}$ and φ_i, φ_{ij} be the covariant derivatives of φ with respect to the metric $g_{\mathbb{S}^{n-1}}$. Define

$$v = \sqrt{1 + |\nabla \varphi|^2_{\mathbb{S}^{n-1}}}.$$

From [19], we know that

$$\lambda = O(e^{\frac{t}{n-1}}), \quad |\nabla \varphi|_{\mathbb{S}^{n-1}} + |\nabla^2 \varphi|_{\mathbb{S}^{n-1}} = O(e^{-\frac{t}{n-1}}). \tag{14}$$

Since $\lambda' = \sqrt{1 + \lambda^2}$, we have

$$\lambda' = \lambda \left(1 + \frac{1}{2} \lambda^{-2} + O(e^{-\frac{4t}{n-1}}) \right). \tag{15}$$

From (14), we also have

$$\frac{1}{v} = 1 - \frac{1}{2}|\nabla\varphi|^2_{\mathbb{S}^{n-1}} + O(e^{-\frac{4t}{n-1}}). \tag{16}$$

In terms of φ, we can express the metric and the second fundamental form of Σ as follows,

$$\begin{aligned} g_{ij} =& \lambda^2(\sigma_{ij} + \varphi_i\varphi_j), \\ h_{ij} =& \frac{\lambda'}{v\lambda} g_{ij} - \frac{\lambda}{v}\varphi_{ij}, \end{aligned}$$

where $\sigma_{ij} = g_{\mathbb{S}^{n-1}}(\partial_{\theta^i}, \partial_{\theta^j})$. Denote $a_i = \sum_k \sigma^{ik}\varphi_{ki}$ and note that $\sum_i a_i = \Delta_{\mathbb{S}^{n-1}}\varphi$. By (14), the principal curvatures of Σ_t has the following form

$$\kappa_i = \frac{\lambda'}{v\lambda} - \frac{a_i}{v\lambda} + O(e^{-\frac{4t}{n-1}}), \quad i = 1, \cdots, n-1.$$

Then we have

$$p_1 = \frac{\lambda'}{v\lambda} - \frac{\Delta_{\mathbb{S}^{n-1}}\varphi}{(n-1)v\lambda} + O(e^{-\frac{4t}{n-1}}).$$

By using (15) and (16), we get

$$p_1 = 1 + \frac{1}{2\lambda^2} - \frac{|\nabla\varphi|^2_{\mathbb{S}^{n-1}}}{2} - \frac{\Delta_{\mathbb{S}^{n-1}}\varphi}{(n-1)\lambda} + O(e^{-\frac{4t}{n-1}}).$$

and

$$p_1^2 - 1 = \frac{1}{\lambda^2} - |\nabla\varphi|^2_{\mathbb{S}^{n-1}} - \frac{2}{n-1}\frac{\Delta_{\mathbb{S}^{n-1}}\varphi}{\lambda} + O(e^{-\frac{4t}{n-1}}).$$

On the other hand,

$$\sqrt{\det g} = \left[\lambda^{n-1} + O(e^{\frac{(n-3)t}{n-1}})\right]\sqrt{\det g_{\mathbb{S}^{n-1}}}.$$

So we have

$$\begin{aligned} \int_{\Sigma_t}(p_1^2 - 1)d\mu =& \int_{\mathbb{S}^{n-1}} \lambda^{n-1}(p_1^2 - 1)dvol_{\mathbb{S}^{n-1}} + O(e^{\frac{(n-5)t}{n-1}}) \\ =& \int_{\mathbb{S}^{n-1}} \left(\lambda^{n-3} - \lambda^{n-1}|\nabla\varphi|^2_{\mathbb{S}^{n-1}}\right) dvol_{\mathbb{S}^{n-1}} \\ & - \frac{2}{n-1}\int_\Sigma \lambda^{n-2}\Delta_{\mathbb{S}^{n-1}}\varphi dvol_{\mathbb{S}^{n-1}} + O(e^{\frac{(n-5)t}{n-1}}) \\ =& \int_{\mathbb{S}^{n-1}} \left(\lambda^{n-3} - \lambda^{n-1}|\nabla\varphi|^2_{\mathbb{S}^{n-1}}\right) dvol_{\mathbb{S}^{n-1}} \\ & + \frac{2(n-2)}{n-1}\int_\Sigma \lambda^{n-3}\langle\nabla\lambda, \nabla\varphi\rangle_{\mathbb{S}^{n-1}} dvol_{\mathbb{S}^{n-1}} + O(e^{\frac{(n-5)t}{n-1}}). \end{aligned}$$

Since $\nabla\lambda = \lambda\lambda'\nabla\varphi$, it follows that $|\nabla\lambda - \lambda^2\nabla\varphi|_{g_{\mathbb{S}^{n-1}}} \leq O(e^{-\frac{t}{n-1}})$, we deduce that

$$\int_{\Sigma_t}(p_1^2 - 1)d\mu = \int_{\mathbb{S}^{n-1}} \left(\lambda^{n-3} + \frac{n-3}{n-1}\lambda^{n-5}|\nabla\lambda|^2\right) dvol_{\mathbb{S}^{n-1}} + O(e^{\frac{(n-5)t}{n-1}}). \tag{17}$$

Moreover,

$$|\Sigma_t|^{\frac{n-3}{n-1}} = \left(\int_{\mathbb{S}^{n-1}} \lambda^{n-1} dvol_{\mathbb{S}^{n-1}}\right)^{\frac{n-3}{n-1}} + O(e^{\frac{(n-5)t}{n-1}}).$$

Using Lemma 4, we achieve

$$\liminf_{t\to\infty} |\Sigma_t|^{-\frac{n-3}{n-1}} \int_{\Sigma_t} (p_1^2 - 1)d\mu \geq \omega_{n-1}^{\frac{2}{n-1}}.$$

□

4. Monotonicity

In this section, we show that $Q(t)$ is monotone non-increasing under IMCF (8).

Proposition 6. *Under IMCF (8), the quantity $Q(t)$ is monotone non-increasing. Moreover, $\frac{d}{dt}Q(t) = 0$ at some time t if and only if Σ_t is totally umbilical.*

Proof. Under IMCF (8), we have

$$\begin{aligned}
&\frac{d}{dt}\int_{\Sigma_t} (p_1^2 - 1)d\mu \\
=&\int_{\Sigma_t} 2p_1\partial_t p_1 d\mu + \int_{\Sigma_t} (p_1^2 - 1)d\mu \\
=&-\frac{2}{(n-1)^2}\int_{\Sigma_t} p_1\left[\Delta\left(\frac{1}{p_1}\right) + \frac{1}{p_1}(|A|^2 - (n-1))\right]d\mu + \int_{\Sigma_t} (p_1^2 - 1)d\mu \\
=&-\frac{2}{(n-1)^2}\int_{\Sigma_t} \left[\frac{1}{p_1^2}|\nabla p_1|^2 + (|A|^2 - (n-1))\right]d\mu + \int_{\Sigma_t} (p_1^2 - 1)d\mu \\
\leq&\frac{n-3}{n-1}\int_{\Sigma_t} (p_1^2 - 1)d\mu,
\end{aligned}$$

where the last inequality follows from the trace inequality $|A|^2 \geq (n-1)p_1^2$. Combining with Proposition 5, we know that the quantity

$$\int_{\Sigma_t} (p_1^2 - 1)d\mu$$

is positive under IMCF (8). Therefore, combining with (10) we get

$$\frac{d}{dt}Q(t) \leq 0.$$

If the equality holds, then the trace inequality implies that Σ_t is totally umbilical. □

Now we finish the proof of Theorem 1.

Proof of Theorem 1. Since $Q(t)$ is monotone non-increasing, we have

$$Q(0) \geq \liminf_{t\to\infty} Q(t) \geq \omega_{n-1}^{\frac{2}{n-1}}. \tag{18}$$

This implies that $\Sigma_0 = \Sigma$ satisfies

$$\int_{\Sigma} (p_1^2 - 1)d\mu \geq \omega_{n-1}^{\frac{2}{n-1}}|\Sigma|^{\frac{n-3}{n-1}},$$

which is equivalent to

$$\int_{\Sigma} p_1^2 d\mu \geq \omega_{n-1}^{\frac{2}{n-1}}|\Sigma|^{\frac{n-3}{n-1}} + |\Sigma|.$$

Now if we assume that equality in (7) is attained, then $Q(t)$ is a constant. Then Proposition 6 indicates that Σ_t is totally umbilical and therefore a geodesic sphere.

If Σ is a geodesic sphere of radius r, then $|\Sigma| = \omega_{n-1}\sinh^{n-1} r$ and $p_1 = \coth r$. Hence, we have

$$\int_\Sigma p_1^2 d\mu = \omega_{n-1}\sinh^{n-1} r\coth^2 r = \omega_{n-1}^{\frac{2}{n-1}}|\Sigma|^{\frac{n-3}{n-1}} + |\Sigma|.$$

Therefore, equality in (7) holds on a geodesic sphere. This finishes the proof of Theorem 1. □

References

[1] A.D. Alexandrov: Zur Theorie der gemischten Volumnia von konvexen Körpern, II: Neue Ungleichungen zwischen den gemischten Volumina und ihre Anwendungen, *Mat. Sb.* **44**(1937), 1205-1238.

[2] A.D. Alexandrov: Zur Theorie der gemischten Volumnia von konvexen Körpern, III: Die Erweiterung zweeier Lehrsatze Minkowskis über die konvexen Polyeder auf beliebige konvexe Flachen, *Mat. Sb.* **45**(1938), 27-46.

[3] W. Beckner: Sharp Sobolev inequalities on the sphere and the Moser-Trudinger inequality, *Ann. of Math.* **138**(1993), 213-242.

[4] A.A. Borisenko, V. Miquel: Total curvatures of convex hypersurfaces in hyperbolic space, *Illinois J. Math.* **43**(1999), 61-78.

[5] S. Brendle: Constant mean curvature surfaces in warped product manifolds, *Publ. Math. Inst. Hautes Études Sci.* **117**(2013), 247-269.

[6] S. Brendle, P.-K. Hung, M.-T. Wang: A Minkowski inequality for hypersurfaces in the Anti-de Sitter-Schwarzschild manifold, *Comm. Pure Appl. Math.* **69**(2016), 124-144.

[7] S.-Y. A. Chang, Y. Wang: On Aleksandrov-Fenchel inequalities for k-convex domains, *Milan J. Math.* **79**(2011), 13-38.

[8] S.-Y. A. Chang, Y. Wang: Inequalities for quermassintegrals on k-convex domains, *Adv. Math.* **248**(2013), 335C377.

[9] S.Y. A Chang, Y. Wang: Some higher order isoperimetric inequalities via the method of optimal transport, *Int. Math. Res. Not.* **24**(2014), 6619-6644.

[10] B.Y. Chen: On total curvature of immersed manifolds, I, *Amer. J. Math.* **93**(1971), 148-162.

[11] B.Y. Chen: Some conformal inariants of submanifolds and their applications, *Boll. Un. Math. Ital.*, **10**(1974), 380-385.

[12] X. Cheng, D. Zhou: Rigidity for nearly umbilical hypersurfaces in space forms, preprint, 2012, arXiv:1208.1786.

[13] L. de Lima, F. Girão: An Alexandrov-Fenchel-Type inequality in hyperbolic space with an application to a Penrose inequality, Ann. Henri Poincaré **17**(2016), 979-1002.

[14] Q. Ding: The inverse mean curvature flow in rotationally symmetric spaces, *Chinese Annals of Mathematics*, Series B, 1-18 (2010).

[15] W. Fenchel: Inégalités quadratiques entre les olumes mixtes des corps convexes, *C. R. Acad. Sci., Paris* **203**(1936), 647-650.

[16] E. Gallego, G. Solanes: Integral geometry and geometric inequalities in hyperbolic space, *Differential Geom. Appl.* **22**(2005), 315-325.

[17] Y.-X. Ge, G.-F. Wang, J. Wu: Hyperbolic Alexandrov-Fenchel quermassintegral inequalities I, preprint, 2013, arXiv: 1303.1714.

[18] Y.-X. Ge, G.-F. Wang, J. Wu: Hyperbolic Alexandrov-Fenchel quermassintegral inequalities II, *J. Differential Geom.* **98**(2014), 237-260.

[19] C. Gerhardt: Inverse curvature flows in hyperbolic space, *J. Differential Geom.* **89**(2011), 487-527.

[20] G.M. Gibbons: Collapsing shells and the isoperimetric inequality for black holes, *Classical Quantum Gravity* **14**(1997), 2905-2915.

[21] P. Guan, J. Li: The quermassintegral inequalities for k-convex starshaped domains, *Adv. Math.* **221**(2009), 1725-1732.

[22] Huisken G., Ilmanen T.: The Riemannian Penrose inequality, *Int. Math. Res. Not.* **20**(1997), 1045-1058.

[23] Huisken G., Ilmanen T.: The inverse mean curvature flow and the Riemannian Penrose inequality, *J. Differential Geom.* **59**(2001), 353-437.

[24] P.-K. Hung, M.-T. Wang: Inverse mean curvature flows in the hyperbolic 3-space revisited, *Calc. Var. Partial Differential Equations* **54**(2015), 119-126.
[25] H.-Z. Li, Y. Wei, C.-W. Xiong: A geometric inequality on hypersurface in hyperbolic space, *Adv. Math.* **253**(2014), 152-162.
[26] P. Li, S.T. Yau: A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces, *Invent. Math.* **69**(1982), 269-291.
[27] M. Maeda: The integral of the mean curvature, *Sci. Rep. Yokohama Nat. Univ.* **25**(1978), 17-21.
[28] G.H. Qiu: A family of higher-order isoperimetric inequalities, *Commun. Contemp. Math.* **17**(2015), 1450015.
[29] M. Ritoré: Optimal isoperimetric inequalities for three-dimensional Cartan-Hadamard manifolds, *Global theory of minimal surfaces*, Clay. Math. Proc., vol. 2, Amer. Math. Soc. Providence, RI, 2005, pp. 395-404.
[30] P. Topping: Mean curvature flow and geometric inequalities, *J. Reine Angew. Math.* **503**(1998), 47-61.
[31] N. Trudinger: Isoperimetric inequalities for quermassintegrals, *Ann. Inst. H. Poincarè Anal. Non Linèaire* **11**(1994), 411-425.
[32] G.-F. Wang, C. Xia: Isoperimetric type problems and Alexandrov-Fenchel type inequalities in the hyperbolic space, *Adv. Math.* **259**(2014), 532-556.
[33] K. Wang: Singularities of mean curvature flow and isoperimetric inequalities in $\mathbb{H}^3$, *Proc. Amer. Math. Soc.* **142**(2015), 2651-2660.

Center of Mathematical Sciences, Zhejiang University, Hangzhou 310027, China,
E-mail address: huyingxiang10@163.com